

# CATEGORIES, ABELIAN CATEGORIES AND COHOMOLOGIES.

YOSHIFUMI TSUCHIMOTO

Examples of derived functors(2)

 Tensor products and Tor.

DEFINITION 9.1. Let  $A$  be an associative unital (but not necessarily commutative) ring. Let  $L$  be a right  $A$ -module. Let  $M$  be a left  $A$ -module. For any  $(\mathbb{Z}$ -)module  $N$ , an map

$$\varphi : L \times M \rightarrow N$$

is called an  **$A$ -balanced biadditive map** if

- (1)  $\varphi(x_1 + x_2, y) = \varphi(x_1, y) + \varphi(x_2, y) \quad (\forall x_1, \forall x_2 \in L, \forall y \in M)$ .
- (2)  $\varphi(x, y_1 + y_2) = \varphi(x, y_1) + \varphi(x, y_2) \quad (\forall x \in L, \forall y_1, \forall y_2 \in M)$ .
- (3)  $\varphi(xa, y) = \varphi(x, ay) \quad (\forall x \in L, \forall y \in M, \forall a \in A)$ .

PROPOSITION 9.2. *Let  $A$  be an associative unital (but not necessarily commutative) ring. Then for any right  $A$ -module  $L$  and for any left  $A$ -module  $M$ , there exists a  $(\mathbb{Z}$ -)module  $X_{L,M}$  together with a  $A$ -balanced map*

$$\varphi_0 : L \times M \rightarrow X_{L,M}$$

*which is universal among  $A$ -balanced maps.*

DEFINITION 9.3. We employ the assumption of the proposition above. By a standard argument on universal objects, we see that such object is unique up to a unique isomorphism. We call it the **tensor product** of  $L$  and  $M$  and denote it by

$$L \otimes_A M.$$

- LEMMA 9.4. (1)  $A \otimes_A M \cong M$ .  
 (2)  $(L_1 \oplus L_2) \otimes_A M \cong (L_1 \otimes_A M) \oplus (L_2 \otimes_A M)$ .  
 (3) For any  $M$ ,  $L \mapsto L \otimes_A M$  is a right exact functor.

DEFINITION 9.5. For any left  $A$ -module  $M$ , the left derived functor  $L_j F(M)$  of  $F_M = \bullet \otimes_A M$  is called the Tor functor and denoted by  $\text{Tor}_j^A(\bullet, M)$ .

By definition,  $\text{Tor}_j^A(L, M)$  may be computed by using projective resolutions of  $L$ .

DEFINITION 9.6. For any group  $G$ , the derived functor of a functor

$$F_G : (G - \text{modules}) \rightarrow (\text{modules})$$

defined by

$$M \mapsto M_G = M / (\mathbb{Z} - \text{span}\{g.m - m; g \in G, M \in M\})$$

is called the homology of  $G$  with coefficients in  $M$ . We denote the homology group  $L_j F_G(M)$  by  $H_j(G; M)$ .

LEMMA 9.7.

$$H_j(G; M) \cong \text{Tor}_j^{\mathbb{Z}[G]}(\mathbb{Z}, M)$$