

$\mathbb{Z}_p, \mathbb{Q}_p$, AND THE RING OF WITT VECTORS

No.12: The ring of Witt vectors and \mathbb{Z}_p

DEFINITION 12.1. Let A be a ring of characteristic p . We call the ring

$$\mathcal{W}_n^{(p)}(A) = \mathcal{W}^{(p)}(A)/V_p^n(\mathcal{W}^{(p)}(A))$$

the ring of Witt vectors of length n . Its elements are called **Witt vectors of length n .**

Note that

$$\mathcal{W}^{(p)}(A)/V_p^n(\mathcal{W}^{(p)}(A))$$

may be considered as a set A^n with an unusual ring structure.

PROPOSITION 12.2.

$$\mathbb{Z}_p \cong \mathcal{W}^{(p)}(\mathbb{F}_p).$$

PROOF. Since $\mathcal{W}^{(p)}$ is a unital commutative ring, there naturally exists a natural ring homomorphism

$$\iota : \mathbb{Z} \rightarrow \mathcal{W}^{(p)}(\mathbb{F}_p).$$

Let us first fix a positive integer n and examine the kernel K_n of a map

$$\pi_n \circ \iota : \mathbb{Z} \rightarrow \mathcal{W}^{(p)}(\mathbb{F}_p)/V_p^n(\mathcal{W}^{(p)}(\mathbb{F}_p))$$

where π_n is the natural projection. Since

$$\#\mathcal{W}^{(p)}(\mathbb{F}_p)/V_p^n(\mathcal{W}^{(p)}(\mathbb{F}_p)) = \#(\mathbb{F}_p^n) = p^n,$$

we have

$$\#(\mathbb{Z}/K_n) | p^n.$$

In other words, $K_n = p^s \mathbb{Z}$ for some integer s . On the other hand, we have

$$\pi_n \circ \iota(p^s) = (1 - T)^{p^s} = (1 - T^{p^s})$$

thus

$$\pi_n \circ \iota(p^s) \in K_n \iff s \geq n.$$

This implies that $K_n = p^n \mathbb{Z}$ and therefore we have an inclusion

$$\mathbb{Z}/p^n \mathbb{Z} \hookrightarrow \mathcal{W}_n^{(p)}(\mathbb{F}_p)$$

which turns to be a bijection ($\because \#(\mathbb{Z}/p^n \mathbb{Z}) = \#(\mathcal{W}_n^{(p)}(\mathbb{F}_p))$).

We then take a projective limit of the both hand sides and obtain the desired isomorphism.

□