

# $\mathbb{Z}_p, \mathbb{Q}_p$ , AND THE RING OF WITT VECTORS

YOSHIFUMI TSUCHIMOTO

No.02: definition of  $\mathbb{Z}_p$

Let  $p$  be a prime (“base”). We would like to introduce a metric on  $\mathbb{Z}$  such that

$$n \text{ :small} \iff n \text{ is divisible by powers of } p.$$

Namely:

DEFINITION 2.1. Let  $p$  be a prime number.

(1) We define a  **$p$ -adic norm**  $|\bullet|_p$  on  $\mathbb{Z}$  as follows.

$$|n|_p = \begin{cases} \frac{1}{p^k} & \text{if } n \neq 0 \text{ and } p^k | n \text{ and } p^{k+1} \nmid n \\ 0 & \text{if } n = 0 \end{cases}$$

(2) We define a  **$p$ -adic distance**  $d_p(\bullet, \bullet)$  on  $\mathbb{Z}$  as follows.

$$d_p(n, m) = |n - m|_p \quad (n, m \in \mathbb{Z})$$

LEMMA 2.2. (1)  $|\bullet|_d$  enjoys the following properties.

- (a)  $|x|_p \geq 0 \quad (\forall x \in \mathbb{Z}). \quad |x|_p = 0 \iff x = 0.$
- (b)  $|x + y|_p \leq \max(|x|_p, |y|_p) \quad (\leq (|x|_p + |y|_p)).$
- (c)  $|xy|_p = |x|_p |y|_p$

(2)  $(\mathbb{Z}, d_p)$  is a metric space.

DEFINITION 2.3. A metric space  $(X, d)$  is said to be **complete** if every Cauchy sequence of  $X$  converges to an element of  $X$ .

THEOREM 2.4. Let  $(X, d)$  be a metric space. There exists a complete metric space  $(\bar{X}, d)$  with an isometry  $\iota : X \rightarrow \bar{X}$  such that  $X$  is dense in  $\bar{X}$ . Furthermore,  $\bar{X}$  is unique up to a unique isometry.

DEFINITION 2.5. Let  $(X, d)$  be a metric space. We call  $(\bar{X}, d)$  as in the above theorem **the completion** of  $(X, d)$ .

DEFINITION 2.6. Let  $p$  be a prime number. We denote the completion of  $(\mathbb{Z}, d_p)$  by  $(\mathbb{Z}_p, d_p)$  and call it **the ring of  $p$ -adic integers**. Thus elements of  $\mathbb{Z}_p$  are  **$p$ -adic integers**.

THEOREM 2.7.  $\mathbb{Z}_p$  has a unique structure of a topological ring. Namely,

(1) There exist unique continuous maps

$$+ : \mathbb{Z}_p \times \mathbb{Z}_p \rightarrow \mathbb{Z}_p$$

(addition) and

$$\times : \mathbb{Z}_p \times \mathbb{Z}_p \rightarrow \mathbb{Z}_p$$

(multiplication) which are extensions of the usual addition and multiplication of  $\mathbb{Z}$ .

(2)  $(\mathbb{Z}_p, +, \times)$  is a commutative associative ring.

DEFINITION 2.8. Let  $p$  be a prime number. For any sequence  $\{a_j\}_{j=0}^{\infty}$  such that  $a_j \in \{0, 1, 2, 3, \dots, p-1\}$ , we consider a sequence  $\{s_n\}$  defined by

$$s_n = \sum_{j=0}^n a_j p^j.$$

Then the sequence  $\{s_n\}$  is a Cauchy sequence in  $\mathbb{Z}_p$ . We denote the limit of the sequence as

$$[0.a_0a_1a_2a_3\dots]_p.$$

EXERCISE 2.1. compute

$$[0.1]_3 + [0.2222\dots]_3$$